

**MATHEMATICS 211**

ASSIGNMENT 3

Due: September 24, 2014

01° Let  $J$  be an open interval in  $\mathbf{R}$ . Let  $f$ ,  $g$ , and  $h$  be differentiable functions defined on  $J$  with values in  $\mathbf{R}$ , for which:

$$\begin{pmatrix} f(t) \\ g(t) \\ h(t) \end{pmatrix} \times \begin{pmatrix} f'(t) \\ g'(t) \\ h'(t) \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad (t \in J)$$

Let  $\Gamma$  be the corresponding mapping carrying  $J$  to  $\mathbf{R}^3$ , with components  $f$ ,  $g$ , and  $h$ :

$$\Gamma(t) \equiv \begin{pmatrix} f(t) \\ g(t) \\ h(t) \end{pmatrix} \quad (t \in J)$$

Let  $\Gamma$  satisfy the Equation of Newton:

$$\Gamma''(t) \equiv \begin{pmatrix} f''(t) \\ g''(t) \\ h''(t) \end{pmatrix} = -\frac{1}{\|\Gamma(t)\|^3} \Gamma(t) \quad (t \in J)$$

Show that the range of  $\Gamma$  is included in a plane. To that end, compute the derivative of:

$$\Delta(t) = \Gamma'(t) \times \Gamma(t)$$

02° Let  $X$  and  $Y$  be nonempty closed subsets of  $\mathbf{R}^2$  for which  $X \cap Y = \emptyset$ . Let  $d$  be the *distance* between  $X$  and  $Y$ , defined as follows:

$$d = \inf\{\|x - y\| : x \in X, y \in Y\}$$

Show by example that  $d$  may be 0 (even though  $X$  and  $Y$  have no point(s) in common). For contrast, show that if  $X$  or  $Y$  is compact then  $d$  is in fact positive.

03° Let  $P$  be the subset of  $\mathbf{R}^2$  consisting of all positions:

$$\begin{pmatrix} x \\ y \end{pmatrix}$$

for which  $y = x^2$ . Let  $\tau$  be the position:

$$\tau = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

in  $\mathbf{R}^2$ . Find the distance between  $P$  and  $\{\tau\}$ .

04• Let  $\mathbf{S}^2$  be the *unit sphere* in  $\mathbf{R}^3$ , consisting of all positions:

$$x = (x_1, x_2, x_3)$$

for which:

$$x_1^2 + x_2^2 + x_3^2 = 1$$

Let  $a, b, c$ , and  $d$  be any four positions:

$$a = (a_1, a_2, a_3)$$

$$b = (b_1, b_2, b_3)$$

$$c = (c_1, c_2, c_3)$$

$$d = (d_1, d_2, d_3)$$

in  $\mathbf{S}^2$  for which:

$$(*) \quad a \neq b, a \neq c, a \neq d, b \neq c, b \neq d, c \neq d$$

Let  $f$  be the function of the foregoing four positions, defined as follows:

$$f(a, b, c, d) = \frac{1}{\|a - b\|} + \frac{1}{\|a - c\|} + \frac{1}{\|a - d\|} + \frac{1}{\|b - c\|} + \frac{1}{\|b - d\|} + \frac{1}{\|c - d\|}$$

Of course, the domain of  $f$  would be the subset  $\Sigma$  of:

$$\mathbf{R}^3 \times \mathbf{R}^3 \times \mathbf{R}^3 \times \mathbf{R}^3 = \mathbf{R}^{12}$$

consisting of all quadruples:

$$(a, b, c, d)$$

of positions in  $\mathbf{S}^2$  which satisfy condition (\*). Show that the range of  $f$  has the form:

$$[\ell, \rightarrow)$$

where  $\ell$  is a suitable positive (!) real number. We mean to say that  $\ell$  is the minimum value of  $f$  but that the values of  $f$  are arbitrarily large. Guess the form of the various quadruples  $(a, b, c, d)$  in  $\Sigma$  for which:

$$f(a, b, c, d) = \ell$$

Start by guessing the “shape” of such quadruples.

05• Reduce the foregoing problem to pairs and triple of positions in  $\mathbf{S}^2$ . Then generalize the foregoing problem to  $k$ -tuples of positions in  $\mathbf{S}^2$ , where  $k$  is any positive integer ( $2 \leq k$ ).