

MATHEMATICS 331

ASSIGNMENT 9

Due: April 9, 2015

01° On \mathbf{R}^3 , we have the conventional pdo geometry, defined by the following bilinear form:

$$\langle\langle \mathbf{x}, \mathbf{y} \rangle\rangle = x_1y_1 + x_2y_2 + x_3y_3$$

where \mathbf{x} and \mathbf{y} are any members of \mathbf{R}^3 :

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

Convert the basis:

$$B_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad B_2 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad B_3 = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$$

for \mathbf{R}^3 to an orthonormal basis, causing minimal disturbance.

02° Let \mathbf{V} be the linear space consisting of all polynomials h with real coefficients, having degree no greater than 3:

$$h(x) = c_0 + c_1x + c_2x^2 + c_3x^3$$

Let \mathbf{V} be supplied with a pdo geometry, as follows:

$$\langle\langle f, g \rangle\rangle = \int_{-1}^1 f(x)g(x)dx$$

where f and g are any polynomials in \mathbf{V} . Introduce the following basis:

$$\mathcal{B}: \quad b_0, b_1, b_2, b_3$$

for \mathbf{V} , where:

$$b_j(x) = x^j \quad (0 \leq j \leq 3, x \in \mathbf{R})$$

Convert \mathcal{B} to an orthonormal basis for \mathbf{V} , causing minimal disturbance. Now let S be the linear mapping in $\mathbf{L}(\mathbf{V})$, defined by differentiation:

$$S(h) = h'$$

where h is any polynomial in \mathbf{V} . Describe the adjoint T of S . Is S self adjoint?

03° Let \mathbf{V}' and \mathbf{V}'' be pdo geometries. Let S and T be a linear mappings in $\mathbf{L}(\mathbf{V}', \mathbf{V}'')$ and $\mathbf{L}(\mathbf{V}'', \mathbf{V}')$, respectively. Show that if S and T are adjoints of one another then the compositions TS and ST in $\mathbf{L}(\mathbf{V}')$ and $\mathbf{L}(\mathbf{V}'')$, respectively, are self adjoint.

04° Let \mathbf{V} be a pdo geometry. Let P be a linear mapping in $\mathbf{L}(\mathbf{V})$ for which $PP = P$. Show that the conditions:

- (1) $\mathbf{V} = \text{ran}(P) \perp \text{ker}(P)$
- (2) P is self adjoint

are equivalent. Under the condition $PP = P$, we say that P is a *projection*. Sometimes, we say “orthogonal projection” rather than “self adjoint projection.” Take special note of condition (1). It figures in both the Spectral Theorem and the Singular Value Decomposition. Verify that if P is a self adjoint projection then $Q = I - P$ is also a self adjoint projection, while:

$$\text{ran}(Q) = \text{ker}(P), \quad \text{ker}(Q) = \text{ran}(P)$$