

The Retinex Theory of Color Vision

A retina-and-cortex system (retinex) may treat a color as a code for a three-part report from the retina, independent of the flux of radiant energy but correlated with the reflectance of objects

by Edwin H. Land

The scientific tradition of simplifying the conditions of an experiment has left us until recently without a satisfactory explanation of how the eye sees color in everyday life. Paradoxically the modern technology of color photography has reinforced the belief that the colors discerned by Newton in the spectrum are, with minor qualifications, the colors of the world around us. We know, for example, that if we use daylight color film when we take a picture in the light shed by an ordinary tungsten-filament lamp, the picture will turn out to have a strong reddish cast. That, we say, is because the rays from the tungsten filament are too "red," never asking how we ourselves can move constantly in and out of tungsten-lit worlds without experiencing any change in the color of familiar objects: apples, lemons, strawberries, bread, human faces (the tones of which are so hard to get right on a television screen).

How, then, does the eye deal with the excess of "red" in a tungsten-lit room? As I hope to demonstrate in this article, the eye, in determining color, never perceives the extra red because it does not depend on the flux of radiant energy reaching it. The eye has evolved to see the world in unchanging colors, regardless of always unpredictable, shifting and uneven illumination. How the eye achieves this remarkable feat has fascinated me for many years.

In 1959 I described in these pages a series of experiments in which a scene created by the superposition of two black-and-white transparencies, one projected through a red filter and the other projected without a filter (that is, in white light), conveys to the eye nearly the gamut of colors present in the original scene [see "Experiments in Color Vision," by Edwin H. Land; *SCIENTIFIC AMERICAN*, May, 1959]. To produce such "red-and-white" images the picture projected through the red filter is taken

through a red filter and the picture projected in white light is taken through a green filter. It would be expected that the superposed image on the projection screen could generate only red, white and various shades of pink. Actually one sees a picture remarkably similar to the full-color photograph reproduced on the opposite page. In the red-and-white photographic projection peppers are green, radishes and strawberries are red, the orange is orange, the lemon and bananas are pale yellow, the wood cutting board and knife handle are brown and the design on the plate is blue.

The challenge presented by our early red-and-white experiments led us step by step over a 20-year period to an explanation of how the visual system is able to extract reliable color information from the world around us, a world in which virtually every scene is lighted unevenly, in which the spectral composition of the radiation falling on a scene can vary enormously and in which illumination as brief as a lightning flash suffices for the accurate identification of color. If the nature of the responses of the photoreceptors in the retina of the eye even approximated what most of us were taught in school, functioning primarily as intensity-level meters with peaks in three different parts of the spectrum, we would be continually confusing one color with another. An object that looked yellow in one part of our field of view might look green or gray or even red when moved to a different part of the field. The fact remains that objects retain their color identity under a great variety of lighting conditions. This constancy is not a minor second-order effect but is so fundamental as to call for a new description of how we see color.

The visual pigments are photosensitive molecules that respond to a wide band of light frequencies. The three pigments in the cone cells of the

retina cover the visible spectrum in three broad, overlapping curves. The pigment with a peak sensitivity at a wavelength of 440 nanometers responds in some degree to the entire lower-frequency half of the visible spectrum. Each of the other two pigments responds to almost two-thirds of the visible spectrum, the two being offset at their peaks by barely 30 nanometers, with their peak sensitivities located at 535 and 565 nanometers [see upper illustration on page 110].

In this discussion the names of colors—"red," "green," "blue" and so on—will be reserved for the color sensation we have when we look at the world around us. In short, only our eyes can categorize the color of objects; spectrophotometers cannot. This point is not a trivial one because many people viewing some of our experiments for the first time will identify something as being red or green but will then ask, as if their eyes were being fooled, "What color is it really?" The answer is that the eye is not being fooled. It is functioning exactly as it must with involuntary reliability to see constant colors in a world illuminated by shifting and unpredictable fluxes of radiant energy.

Since I believe the study of color in fully colored images is best begun by examining images that are completely devoid of and completely uncomplicated by the experience of color, let me describe that experience in some detail. The hypersensitive system based on the rod cells in the retina functions at light levels as much as 1,000 times weaker than the systems based on the cone cells do, so that it is possible to answer the interesting question: What colors will one see if only the rod system is activated? One procedure is to put on a pair of tightly fitting goggles equipped with neutral-density filters that reduce the incident light by a factor of 30,000. After one has worn the goggles for about half

an hour objects in a room illuminated to the typical level of 20 foot-candles will become visible. The effective illumination in the room will thus be 1/1,500 foot-candle. As one looks around the room the familiar colored objects will be seen devoid of color, exhibiting instead a range of lightnesses from white to black, much as they would appear

in a black-and-white photograph taken through a green color-separation filter. In other words, the reds will appear very dark, the greens lighter, the blues dark, the whites light and the blacks very dark.




In this colorless world one finds that the nature of the image is not determined by the flux of radiant energy

reaching the eye. The illumination can easily be arranged so that there is more flux from a region that continues to look very dark than there is from a region that continues to look very light, whether these regions are three-dimensional objects or artifacts contrived with a montage of dark and light pieces of paper. The paradox immediately arises



STILL LIFE was used to make the four black-and-white images presented below. The reproduction of the still life above was made by conventional processes of color photography and photoengraving to show the reader what the colors of the original objects in the scene

were. The black-and-white images were made with film-filter combinations that closely duplicate the separate wavelength sensitivities of the four systems of photoreceptors in the retina of the eye: the three systems of cone cells and the hypersensitive system of rod cells.



BLACK-AND-WHITE IMAGES OF STILL LIFE were taken with four different film-filter combinations, creating what the author calls retinex records. The picture at the top left was taken with a film whose spectral response was altered so that it matched the spectral sensitivity of the long-wave cone pigments in the eye. This photograph enables the observer to see a colorless image that approximates the image produced by the long-wave cones by themselves. The pic-

ture at the top right shows the same scene as it would be viewed by the middle-wave cone pigment. The picture at the bottom left is the scene as it would be viewed by the short-wave cone pigment. The picture at bottom right corresponds to the image seen by the rods. Unlike cone images, which cannot be viewed independently, images produced by the rod pigment can be studied in isolation at very low light levels, without interference from much less sensitive cone systems.

that each of the objects, the pieces of paper for example, whether dark or light or in between, maintains its lightness without significant change as it is moved around the room into regions of higher or lower flux. Light papers will be seen as being light and dark papers simultaneously as being dark, even with the same flux coming from each of them to the eye. Strong gradients of flux across the field will be apparent only weakly, if at all.

Furthermore, in an intricate collage of areas of various lightnesses, sizes and shapes, the lightness of a given element does not change visibly as it is relocated

in any part of the collage and associated with a new arbitrary surround. When a small area is totally surrounded by a large area, the lightness of the small area will change somewhat depending on whether the large area is darker or lighter than the small one. In general, however, the impressive fact is that the lightness of a given area is not appreciably modified by the immediately surrounding areas, nor is it modified by the still larger areas surrounding them.


Although I have been describing a colorless world as it is seen by the hypersensitive receptors of rod vision, all

the observations about the stability of lightness values can readily be reproduced with a montage of white, black and gray papers viewed at ordinary light levels. If, for example, a square of matte-surface black paper or, better still, black velvet is placed at one side of such a montage and a square of white paper is placed at the opposite side several feet away, with an assortment of light and dark papers scattered in between, one can place a strong light source close enough to the black square so that it sends more radiant energy to the eye than the white square, remote from the light; yet the black square will continue to look black and the white square white. In fact, with the montage still strongly illuminated from one side either the black square or the white one can be moved to any other part of the montage without a significant change in its appearance.


This remarkable ability of the eye to discover lightness values independent of flux, so convincingly demonstrated when only a single photoreceptor system is operating, is the rock on which a satisfactory description of color vision can be built. The first response of the visual system is for the receptors to absorb the light falling on the retina. Whereas the initial signal produced in the outer segment of the receptor cell is apparently proportional to the light flux absorbed by the visual pigment, the final comprehensive response of the visual system is "lightness," which shows little or no relation to the light flux absorbed by the visual pigment.

The processing of fluxes to generate lightnesses could occur in the retina, or in the cerebral cortex, or partially in both. Since we are uncertain of the location of the mechanisms that mediate these processes, I have coined the term *retinex* (a combination of retina and cortex) to describe the ensemble of biological mechanisms that convert flux into a pattern of lightnesses. I shall therefore use the term throughout this article in referring to these biological mechanisms. I shall also reserve the term *lightness* to mean the sensation produced by a biological system. Although the rods can be stimulated at light intensities below the cone threshold, the cones cannot be stimulated without exciting the rods. For cones we must study the lightness images produced by each individual set of receptors using *retinex* photography, as I shall explain below, or learn the properties of lightness images from model calculations based on spectroradiometric measurements.


Now that we know that at low light levels an isolated receptor system generates an image in terms of lightness that is completely free of color, might it be possible to bring one of the cone systems into operation along with the hypersensitive system, so that only the



NORMALIZED SPECTRAL SENSITIVITIES OF FOUR VISUAL PIGMENTS (solid lines) span the visual spectrum in overlapping curves. Curve that peaks at about 500 nanometers corresponds to sensitivity of rod pigment. Other three curves represent cone pigments. Broken lines show sensitivities of the film-filter combinations that were selected to match the sensitivities of the four retinal pigments and used to make the black-and-white *retinex* records in the illustration at bottom of preceding page. Cone curves are adapted from work of Paul Brown and George Wald of Harvard University. The rod curve is standard scotopic luminosity curve.




THRESHOLD RESPONSES OF RETINAL RECEPTORS vary by large factors. The hypersensitive rod system provides vision at radiance levels about 1,000 times weaker than the light levels needed to activate the cone systems. It has been shown in author's laboratory that multicolored scenes exhibit nearly their normal range of colors when they are viewed at light levels so adjusted that only rod system and one cone system, the long-wave system, are responding.



“COLOR MONDRIAN” EXPERIMENT employs two identical displays of sheets of colored paper mounted on boards four and a half feet square. The colored papers have a matte finish to minimize specular reflection. Each “Mondrian” is illuminated with its own set of three projector illuminators equipped with band-pass filters and independent brightness controls so that the long-wave (“red”), middle-wave (“green”) and short-wave (“blue”) illumination can be mixed in any desired ratio. A telescopic photometer can be pointed at any area to measure the flux, one wave band at a time, coming to the eye

from that area. The photometer reading is projected onto the scale above the two displays. In a typical experiment the illuminators can be adjusted so that the white area in the Mondrian at the left and the green area (or some other area) in the Mondrian at the right are both sending the same triplet of radiant energies to the eye. The actual radiant-energy fluxes cannot be re-created here because of the limitations of color reproduction. Under actual viewing conditions white area continues to look white and green area continues to look green even though the eye is receiving the same flux triplet from both areas.



IDENTICAL ENERGY FLUXES AT THE EYE provide different color sensations in the Mondrian experiments. In this example, with the illuminants from the long-wave, middle-wave and short-wave illuminators adjusted as indicated, an area that looks red continues to look red (*left*), an area that looks blue continues to look blue (*middle*)

and an area that looks green continues to look green (*right*), even though all three are sending to the eye the same triplet of long-, middle- and short-wave energies. The same triplet can be made to come from any other area: if the area is white, it remains white; if the area is gray, it remains gray; if it is yellow, it remains yellow, and so on.

completely colorless system and one other were functioning? This two-receptor experiment has been carried out and provides a powerful confirmation of the ideas derived from all our binary work with red-and-white images and subsequent ternary studies with multicolored displays seen under various illuminants. The experiment, rapidly becoming a classic, was devised by my colleagues John J. McCann and Jeanne L. Benton. McCann and Benton illuminated a

color display with a narrow wave band of light at 550 nanometers. The light level was raised just above the amount needed to make the display visible to the dark-adapted eye, thus ensuring that only the hypersensitive system was operating. They then added a second narrow-band illuminant at 656 nanometers, with its level adjusted so that it was just sufficient to activate the long-wave receptor system but not the middle-wave receptor system. Under these conditions only two


receptor systems, namely the rods and the long-wave cones, were receiving enough light to function.

The resulting image exhibited a remarkable range of color, enabling an observer to assign to each area in the display the same color name it would have if it were illuminated above the cone threshold. The result is reminiscent of the multicolored images produced by the red-and-white system. The demonstration explicitly confirms our early proposition that the lightness information collected at two wave bands by separate receptor systems is not averaged, point by point and area by area, but is kept distinct and is compared. We know that the rod system does not produce a colored image when the image is seen by itself, and we know that the long-wave light alone cannot produce an image with a variety of colors. The combination, however, gives rise to a wide variety of colors, namely reds, yellows, browns, blue-greens, grays and blacks.


What, then, accounts for the color? The emergence of variegated colors can be ascribed to a process operating somewhere along the visual pathway that compares the lightnesses of the separate images on two wave bands, provided by the two independent retinex systems. The two-receptor experiment makes it plausible that when three independent images constituting the lightnesses of the short-, middle- and long-wave sets of receptors are associated to give a full-colored image, it is the comparison of the respective lightnesses, region by region, that determines the color of each region. The reason the color at any point in an image is essentially independent of the ratio of the three fluxes on three wave bands is that color depends only on the lightness in each wave band and lightness is independent of flux.

As we have seen, the spectral sensitivities of the visual pigments overlap broadly. If we illuminated a scene with the entire range of wavelengths to which a single visual pigment is sensitive, we would see a large variety of colors because more than one retinex system would respond. With the help of filters and appropriate film emulsions, however, we can isolate the lightnesses that would ordinarily be incorporated into the sensation of color. We call black-and-white photographs made for this purpose retinex records.

The photographic technique, making use of silver emulsions, performs two functions. First, the system provides spectral sensitivities that are the same as those of the visual pigments. Second, it generates black-and-white pictures for a human observer to examine. It is the human visual system that converts the photographic pattern deposited in silver into lightness. Ideally we should like




COLOR-MATCHING EXPERIMENT uses a simplified Mondrian of 17 color areas (left) and a standard color reference, *The Munsell Book of Color*, which contains 1,150 color “chips” (right). The Mondrian is illuminated with three narrow-band light sources: one at 630 nanometers (long-wave light), one at 530 nanometers (middle-wave light) and one at 450 nanometers (short-wave light). The ratio of the three illuminants can be adjusted so that the triplet of energies reflected to the eye from any chosen area will exactly equal the triplet that previously reached the eye from some other area. In this experiment five areas, gray, red, yellow, blue and green, were selected in sequence to send the same triplet of energies to the eye. In each of the five consecutive parts of this experiment the observer selected from the Munsell book the chips that came closest to matching the 17 areas of the Mondrian. The Munsell book was illuminated throughout the experiment with a constant spectral mixture of three narrow-band lights adjusted at the outset so that the white Munsell chip appeared the “best white.” The experiment was set up so that the observers used one eye for viewing the Mondrian and the other eye for viewing chips. Gray paper with an opening was used to provide chips with a constant surround.






PROPORTIONS OF NARROW-BAND ILLUMINANTS used to light the simplified Mondrian in the Munsell-chip matching experiments were adjusted as is shown by the bars at the top of this illustration so that five different areas of the Mondrian (indicated by arrows) sent to the observer's eye in successive matching trials the same trip-

let of energies: 5.8 flux units of long-wave light, 3.2 flux units of middle-wave light and 1.6 flux units of short-wave light. The illustration below shows the Munsell chips that were selected in the constant illuminant to match the five Mondrian areas (gray, red, yellow, blue and green) that had sent to the eye exactly the same triplet of energies.



MUNSELL CHIPS SELECTED BY OBSERVERS to match the five Mondrian areas that had sent identical triplets of energy to the eye are reproduced. The Munsell book was illuminated with a constant spectral mixture of narrow-band illuminants (bars at top) and the chips were viewed within a constant gray surround. The energy that

was sent to the eye by the selected Munsell chips is shown by the bars at the bottom of the illustration. It is evident that the match between the Mondrian areas and the Munsell chips is not made on the basis of the flux of radiant energy at the eye of the observer. What does cause the two areas to match is described in the illustrations that follow.



FURTHER ANALYSIS OF MATCHING EXPERIMENT begins to identify the basis on which the visual system makes the color match between the Mondrian area and the Munsell chip without regard to the flux each member of the pair sends to the eye. The efficiency with which a given area in the Mondrian reflects light in each of the three wave bands (first column) multiplied by the amount of energy striking that area in each of the wave bands (second column) yields

the energy triplet that reaches the eye (third column). The three columns at the right contain comparable data for the Munsell chips selected as a match for the Mondrian areas. Whereas illustration at bottom of the preceding page shows that the eye does not match colors using a "meter" that measures triplets of energies at the eye, this illustration shows that when a match is made, it is the reflectances of two areas that correspond, as is shown in first and fourth columns.

our observer to examine the black-and-white pattern with only one set of cones, reporting the lightnesses appropriate to that set. At any point in the black-and-white pattern, however, the reflectance is essentially the same throughout the visible spectrum. Therefore with a black-and-white photograph we stimulate all the receptors with the same information, that is, with the energies that would be absorbed by a single visual pigment. If we assume that all the retinex systems process information in an identical manner, we can propose that sending this identical information to several sets of receptors is the same as sending it to only one receptor, thereby enabling us to see what the image would look like if it were possible to isolate it.

On page 109 the reader will see three black-and-white pictures taken through retinex filters that simulate the response of the three cone pigments. The strawberries and radishes, for example, are light on the long-wave record, darker on the middle-wave record and darkest on the short-wave record. Although the orange and lemon are about as dark as the strawberries and radishes on the short-wave record, they are nearly as light on the middle-wave record as they are on the long-wave record. On the printed page the distinctions are subtle. To the eye viewing an actual full-color scene the subtle distinctions provide all the information needed to distinguish countless shades and tints of every color.

After the three lightnesses of an area have been determined by the three retinex systems no further information is necessary to characterize the color of any object in the field of view. Any specific color is a report on a trio of three specific lightnesses. For each trio of lightnesses there is a specific and unique color.

The limitations of color photography make it impossible to show the reader the demonstrations readily accomplished in our laboratory, which dramatically reveal the independence of perceived color from the flux reaching the eye. What the reader would see would be two boards four and a half feet square identically covered with about 100 pieces of paper of various colors and shapes. In order to minimize the role of specular reflectance the papers have matte surfaces and, except for black, have a minimum reflectance of at least 10 percent for any part of the visible spectrum. In these displays, which we call "color Mondrians" (after the Dutch painter to whose work they bear a certain resemblance), the papers are arranged so that each one is surrounded by at least five or six others of different colors [see top illustration on page 111].

Each of the identical Mondrians is illuminated by its own set of three pro-

jectors equipped with sharply cutting band-pass filters (not retinex filters): one at 670 nanometers embracing a band of long waves, one at 540 nanometers embracing a band of middle waves and one at 450 nanometers embracing a band of short waves. The amount of light from each illuminating projector is controlled by a separate variable transformer. In addition the illuminating projectors have synchronized solenoid-activated shutters to control the duration of illumination. There is a telescopic photometer that can be precisely aimed at any region of either Mondrian to measure the amount of radiation reflected from any point and therefore the amount of flux reaching the eye. The output of the photometer is projected on a scale above the Mondrian, where it can be seen by those taking part in the demonstration.

The demonstration begins with the three illuminating projectors turned on the Mondrian on the left; the Mondrian on the right remains dark. The variable transformers are set so that the entire array of papers in the left Mondrian are deeply colored and at the same time the whites are good whites. This setting is not critical. Then, using one projector at a time and hence only one wave band at a time, we measure with the telescopic photometer the energy reaching the eye from some particular area, say a white rectangle. The readings from the white area (in milliwatts per steradian per square meter) are 65 units of long-wave light, 30 units of middle-wave light and five units of short-wave light. We have now established the three energies associated with that sensation of white.

We turn off the three projectors illuminating the color Mondrian on the left. On the right we turn on only the long-wave projector. We select a different area of unknown color and adjust the long-wave light until the long-wave energy coming to the eye from the selected area is the same as the long-wave energy that a moment ago came from the white paper in the Mondrian on the left, 65 units. We turn off the long-wave projector and separately adjust the transformers controlling the middle- and short-wave projectors, one after the other, so that the energies sent to the eye from the selected area are also the same as those that came from the white area on the left. We have not yet turned on all three light sources simultaneously, but we know that when we do so, the triplet of energies reaching the eye from the selected area of still unknown color will be identical with the triplet that had previously produced the sensation white.

When we turn on the three illuminants, we discover that the area in the Mondrian on the right is green. We now illuminate the Mondrian on the left with its illuminants at their original settings

so that both Mondrians can be viewed simultaneously. The white area on the left continues to look white and the green area on the right continues to look green, yet both are sending to the eye the same triplet of energies: 65, 30 and five in the chosen units.

We turn off the illuminants for both Mondrians and select some other area in the left Mondrian and sequentially adjust the energies reaching the eye from it so that they are the same as the energies that originally gave rise to the sensation of white and also gave rise to the sensation of green in the right Mondrian. When we turn on all three projectors illuminating the left Mondrian, we see that this time the selected area is yellow. The triplet of energies reaching our eye is the same one that had previously produced the sensations of white and green. Again, if we wish, the yellow and green can be viewed simultaneously, with yellow on the left and green on the right.

We can continue the demonstration with other areas such as blue, gray, red and so on. It is dramatically demonstrated that the sensation of color is not related to the product of reflectance times illumination, namely energy, although that product appears to be the only information reaching the eye from the various areas in the Mondrians.

In order to demonstrate that the color sensations in these experiments do not involve extensive chromatic adaptation of retinal pigments the projectors are equipped with synchronized shutters so that the Mondrians can be viewed in a brief flash, a tenth of a second or less in duration. Regardless of the brevity of observation the results of the demonstrations are not altered. Thus one can say that neither chromatic adaptation nor eye motion is involved in producing the observed colors. Finally, the very essence of the design of the color Mondrian is to obviate the significance of the shape and size of surrounding areas, of the familiarity of objects and of the memory of color. Curiously, from time to time there is a casual attempt to induce what is called color constancy as an explanation of these demonstrations. Clearly color constancy is only a compact designation of the remarkable competence that is the subject of this article.

The mystery is how we can all agree with precision on the colors we see when there is no obvious physical quantity at a point that will enable us to specify the color of an object. Indeed, one can say the stimulus for the color of a point in an area is not the radiation from that point. The task of psychophysics is to find the nature of the stimulus for that color.


Here let us remember that what the eye does unfaithfully well is to discover

lightness values independent of flux. We saw this to be true for a single receptor system, the rod system, operating alone and for the three cone systems operating collectively when they viewed an array of white, gray and black papers. Let us now illuminate the colored Mondrian array with light from just one of the three projectors, say the projector supplying long-wave light, and observe the effect of increasing and decreasing the flux by a large factor. We observe that the various areas maintain a constant rank order of lightness. If, however, we switch the illumination to a different wave band, say the middle wave band, the lightnesses of many of the areas will change; many of the 100 or so areas

will occupy a different rank order from lightest to darkest. Under the short-wave-band illuminant there will be yet a third rank order. Specifically, a red paper will be seen as being light in the long-wave light, darker in middle-wave light and very dark in short-wave light. A blue paper, on the other hand, will be light in short-wave light and very dark in both middle- and long-wave light. Papers of other colors will exhibit different triplets of lightnesses. When we conducted such experiments nearly 20 years ago, we were led inevitably to the conclusion that the triplets of lightnesses, area by area, provided the set of constancies we needed to serve as the stimuli for color, independent of flux.

It is evident that the lightnesses exhibited by a given piece of colored paper under illuminants of three different wave bands is related to the amount of energy the paper reflects to the eye at different wavelengths. Let us now examine, by means of a particular experiment, how such reflectances can be related step by step to perceived lightnesses and how, in the process, the radiant flux that reaches the eye—the ultimate source of knowledge about lightness—finally becomes irrelevant to the sensation of color.

In our laboratory McCann, Suzanne P. McKee and Thomas H. Taylor made a systematic study of observers' re-



ROLE OF REFLECTANCE and its psychophysical correlate, lightness, in guiding the eye to match Munsell chips with Mondrian areas was examined with the help of retinex filter-photomultiplier combinations that match the spectral sensitivity of the cone pigments. Under each combination of illuminants (*top*) the integrated radiance, or flux, in each retinex wave band of a Mondrian area was compared with the integrated radiance of a sheet of white paper. The ratio of

integrated radiances yields the integrated reflectance of the Mondrian area, expressed here in percent. For the matching Munsell chip a set of ratios was similarly determined (*bottom*). The final step in deriving a physical equivalent of lightness is the scaling, or spacing, of integrated reflectances to be consistent with the spacing of lightness sensations. This transformation is explained in the illustration on the opposite page. The scaled values appear in the column at the right.

sponses to a simplified color Mondrian with areas of 17 different colors. They asked the observers to match the 17 areas one at a time under different illuminants with colored squares of paper that had been selected from a standard color-reference book, *The Munsell Book of Color* and that were viewed under a constant "white" illumination.


The illuminants on the Mondrian were adjusted in five separate matching experiments so that five different areas (gray, red, yellow, blue and green) sent to the eye an identical triplet of radiances. The observer began by selecting a matching Munsell "chip" for each of the 17 areas in the Mondrian when the gray area in the Mondrian sent a particular triplet of energies to the eye. Another set of 17 matching Munsell chips was selected when the same triplet was later sent to the eye by a red area in the Mondrian, and the same was done for yellow, blue and green areas under illuminants that supplied the same triplet of energies.

The illustrations on page 113 show the details of the experiment and the five different Munsell colors the observers selected to match the five areas when each area sent to the eye precisely the same triplet of energies. In spite of the constancy of the energy reaching one eye from the Mondrian, each observer, using the other eye, selected Munsell chips that were gray, red, yellow, blue and green.

The constant illumination used in viewing the Munsell book was a triplet of illuminants at three wavelengths that observers judged to produce the "best" white. The actual triplet of wavelengths reaching the eye from the whitest paper in the Munsell book was 11.5 units of long-wave light, 7.8 units of middle-wave light and 3.3 units of short-wave light. The illuminants supplied energy in narrow bands with peaks at 630 nanometers, 530 nanometers and 450 nanometers. A similar triplet of narrow-band illuminants were mixed in various proportions to illuminate the Mondrian.

At this point the reader might ask: Would not a single gray area exhibit a pronounced change in color if the surrounding papers had reflected light of widely differing spectral composition? Could these changes in color account for the results of the Mondrian experiments? The answer to the questions is that no manipulation of surrounding papers in the Mondrian is capable of making the gray paper match the red, yellow, blue and green Munsell papers selected by the observers in the Mondrian experiment.

McCann, John A. Hall and I have examined the matter further by repeating the Mondrian-Munsell experiment in various ways so that the average spectral composition of the light reaching



SENSATION OF LIGHTNESS is plotted on an equal-interval scale. Observers are shown a sheet of white paper (9) and a sheet of black paper (1) and are then asked to choose a sheet of paper whose shade of gray lies halfway between the two. The selection is the gray labeled 5. Similar selections are made to determine the locations of midpoints between 1 and 5 and between 5 and 9 and so on until the equal-interval scale is filled. The end values 0 and 10 are extrapolations. The curve is then plotted by measuring the reflectances of the various papers selected by the observers. The curve makes it possible to convert values of integrated reflectance into values of scaled integrated reflectance, as is given in illustration on opposite page.

the eye from the Mondrian and its surround remains the same regardless of the spectral composition of the light needed to establish a constant triplet from area to area. We have done this in one case by surrounding the entire Mondrian with brightly colored papers selected in such a way that they exactly offset the average mixture of wave bands from the Mondrian itself and, more dramatically, by cutting the 17 areas of the Mondrian apart and placing them well separated on the backgrounds of offsetting color. Neither arrangement has any significant effect on the Munsell chips chosen to match the various areas of the Mondrian.

Let us return, then, to the search for the stimulus that guides us so accurately to the correct identification of colors. If it is not a flux of radiant energy at the eye from each point in the field of view, what are the physical correlates of the lightnesses of objects on three separate

wave bands, corresponding to the spectral sensitivities of the cone pigments? Can such a precise physical correlate of lightness be demonstrated?

McCann, McKee and Taylor next measured the radiance, or energy at the eye, of the various Mondrian areas and of the matching Munsell chips by using a photomultiplier in conjunction with a version of the retinex filters. Since the retinex-photomultiplier combination integrates the flux of radiant energy over a broad band of wavelengths, the instrument provides a value we call integrated radiance. McCann and his colleagues then obtained the integrated radiances from a large sheet of white paper placed under each of the experimental illuminants that had been used to light the Mondrian in the chip-matching experiments. If the integrated radiance from a Mondrian area is used as the numerator in a fraction and the integrated radiance from the white paper is used as the denominator, one obtains a value for in-


egrated reflectance, which can be expressed as a percent.

The integrated reflectances for the various Munsell chips are determined in the same manner under the constant "white" illumination. This amounts to measuring the percentage of reflectance using detectors with the same spectral sensitivity as the visual pigments. The results show that the Munsell chip chosen by the eye to match a given Mondrian area will have approximately the same three integrated reflectances as the area. For example, the blue area in the Mondrian has a triplet of integrated reflectances (long-, middle- and short-wave) of 27.3, 35.9 and 60.7 percent. The comparable values for the matched Munsell chip are 34.6, 38.5 and 57.1 percent [see illustration on page 116].

Finally, the integrated reflectances are "scaled" so that their equal spacing is consistent with the equal spacing of lightness sensations. The curve for this transformation is shown in the illustration on the preceding page. Using this curve, we see that the blue area in the Mondrian has a triplet of scaled inte-

grated reflectances of 5.8, 6.5 and 8.1, whereas the corresponding values for the matching Munsell chip are 6.4, 6.7 and 7.9. If we study the five areas that successively sent identical triplets of energies to the eye and compare their scaled integrated reflectances with those of their matching Munsell chips, we find that all the values are in excellent agreement. In other words, in the triplets of integrated reflectances we have identified a highly accurate physical correlate of color sensations. The data fall along the 45-degree line that describes the locus of perfect correlation [see illustration below].

We have sought a physical correlate for lightness, and we have found that the scaled integrated reflectances of the five areas that sent identical triplets of fluxes to our eyes are the same as those of the matching Munsell chip. This correlation enables us to use scaled integrated reflectances as a measured lightness equivalent. The problem now shifts to one of how the eye derives the lightness that corresponds to the reflectances of objects in each wave band.




AGREEMENT IN SCALED INTEGRATED REFLECTANCES between Mondrian areas and Munsell chips chosen to match them is summarized for all three wave-band systems. The scaled integrated reflectances of five Mondrian areas and matching Munsell chips were determined as is described in illustration on page 116. In this graph triplets of scaled integrated reflectances of five Mondrian areas that sent identical fluxes to the eye are plotted against scaled integrated reflectances of Munsell chips chosen to match them. Although the dots collectively represent correspondence for all three retinex wave bands, any particular dot denotes the degree of correspondence on one retinex wave band between a Mondrian area and a Munsell chip. Close correspondences show that scaled integrated reflectance is physical correlate of the sensation "lightness," showing precision with which a triplet of lightnesses determines color.

It is one thing to measure a triplet of lightness equivalents using a retinex filter coupled to a photomultiplier; it is quite another for the eye to determine lightnesses in the unevenly lighted world without reference sheets of white paper. I described above the ability of an isolated receptor system—the hypersensitive system of rod vision—to classify objects correctly according to their inherent reflectivity regardless of whether the objects happened to be in a brightly or a dimly lighted region of visual space. The ability of one receptor system to work in this way makes it plausible that the other three systems of normal daytime vision possess the same ability, each system viewing the world through a broad but restricted region of the spectrum, the regions we duplicate with retinex filters. Each system forms a separate lightness image of the world. The images are not mixed but compared. The comparison of lightnesses at each area gives rise to the range of sensations we know as color.

How could the biological system generate a hierarchy and spacing of lightness values given only the flux from each point in a scene and knowing nothing about the pattern of illumination and nothing about the reflectances of objects? The scheme I am about to describe is the most general we have found that surmounts these limitations; its physiological embodiment could take many forms.


Let me begin by pointing out the significance of edges in defining objects or areas in a scene. If a sheet of white paper is lighted strongly from one side, we see no discontinuity in color from one side to the other. Let us now imagine two light detectors positioned to measure the luminance from two different places on the paper. If the illumination is non-uniform, the luminances of the two places will of course be different. As the two detectors are moved closer together the luminances approach the same value and the ratio of the two outputs approaches unity. If, however, the two detectors bridge the boundary between two areas that differ abruptly in reflectance, such as would be the case with even a pale gray square on the white paper, the ratio of the outputs of the two detectors will approach the ratio of the two reflectances. Thus the single procedure of taking the ratio between two adjacent points can both detect an edge and eliminate the effect of nonuniform illumination. If we process the entire image in terms of the ratios of luminances at closely adjacent points, we can generate dimensionless numbers that are independent of the illumination. These numbers give the ratio of reflectances at the edge between adjacent areas; the reflectances themselves are not yet ascertained.

In order to determine reflectances we need to relate all these ratios of reflec-





THE EYE'S METHOD OF DISCOVERING LIGHTNESS in complex images remains to be established. An efficient and physiologically plausible scheme is depicted in this illustration and the one below. The numbers inside the schematic Mondrian represent the long-wave integrated radiances coming from each area of a display that is evenly lighted. The long-wave retinex system independently "measures" the long-wave integrated radiance, point by point, as if it were doing so along an arbitrary pathway (*color*). The flux at each successive closely spaced pair of points is converted into a ratio. This ratio is subjected to a threshold test: any ratio to be regarded as a change must vary from unity by more than some small threshold amount (plus or minus .003 in the computer program). If the ratio does not vary from unity by this amount, it is regarded as being "unchanged" and is set to equal unity. A second threshold-tested ratio along the same pathway is multiplied by the first ratio to give a sequential product that is both

the model's response for that point and the signal sent along to be multiplied by the next ratio. When the path crosses an edge between two lightnesses, there is a sharp change in the threshold-tested ratio and hence a similar change in the sequential product. Here the path is started in the white area, where the flux of radiant energy is 100. By the time the path reaches the brown area at the lower right the product is .18. The retinex system has thus determined that the brown area reflects 18 percent as much long-wave energy as the white area. Any other path ending in the brown area would yield the same result as long as it had been through the white area. By averaging the responses for each area, as computed by many arbitrary paths, the long-wave retinex system arrives at a single reflectance value for each area, which designates perceived lightness. Middle- and short-wave retinex systems compute their own sets of lightness values. Comparison of triplet of lightnesses for each area provides sensation of color.



MORE REALISTIC CASE OF GRADED ILLUMINATION is handled equally well by the sequential-product method to arrive at the same reflectance value of .18 for the brown area at the end of the path, even though here the long-wave retinex system receives as much flux from the middle of the brown area (57) as it does from the middle

of the white area (57). The scheme hence provides a means for arriving at computed reflectance independent of flux and without resort to white cards as standards. Precise values of light flux along pathway in this diagram were derived from a computer program that works with 75 values between every two values printed within Mondrian.



tances in the field of view. Given the ratio of luminances at the edge between a first area and a second one, we multiply it by the ratio of luminances at the edge between the second area and a third. This product of ratios approaches the ratio of reflectances between the first and third areas, regardless of the distribution of illumination. Similarly, we can obtain the ratio of reflectances of any two areas in an image, however remote they are from each other, by multiplying the ratios of all the boundaries between the starting area and the remote area. We can also establish the ratio of the reflectance of any area on the path by tapping off the sequential product reached at that area [see illustrations on preceding page].

We are now coming close to the answer to the question: How can the eye ascertain the reflectance of an area without in effect placing a comparison standard next to the area? The sequential product can be used as a substitute for the placement of two areas adjacent to each other, thus defining a photometric operation feasible for the eye.

The remaining task is to suggest how the eye can discover the area of highest reflectance in the field of view and then decide whether that area is actually white or some other color. In the model we have proposed, sequential products are computed along many arbitrary pathways that wander through the two-dimensional array of energies on the model's "retina." Since the pathways

COLOR "SOLID" shows the location of all perceivable colors, including white and black, in a three-dimensional color space constructed according to the author's retinex theory. The position of a color in this space is determined not by the triplet of energies at a point but by the triplet of lightnesses computed by the eye for each area. The color photograph at the top left shows the location of representative colors throughout the space. The direction of increasing lightness along each axis is shown by the arrows. The three black-and-white photographs of the color solid were taken with retinex filter-film combinations. They show the lightness values of the representative colors as they would be perceived separately by the eye's long-wave (*top*), middle-wave (*middle*) and short-wave (*bottom*) visual pigments. The set of 10 color pictures at the right represents horizontal planes cut through the three-dimensional color space. Each plane is the locus of colors possible with a constant short-wave lightness. For example, the fifth plane from the bottom shows the variety of color sensations from all possible long- and middle-wave lightness values when those values are combined with a short-wave lightness of 5. The colored squares are samples taken from *The Munsell Book of Color*. In general the blank areas on each plane represent regions where colors could be produced only by fluorescent dyes, if they were produced at all.

The people who developed the 4 hour video home system now present some of the ways it can change your life.

Introducing Omnivision IV™ VHS™, the Panasonic four-hour video home player/recorder from the people who developed the four-hour video home system—the engineers of our parent company.

Four hours of video recording. Instead of two.

Four hours of taping when you're not at home.

So when you know you're going to be stuck at the office, you can also know that four hours of video taping will let you enjoy every moment of the big game...even if it runs into overtime or extra innings.



When work rearranges your schedule, rearrange the network's schedule.

Two great shows on at the same time? Watch one, tape the other.

entire movie...even the climax when the Monster eats Yankee Stadium.

Four hours of video taping instead of two means there's virtually no program, no special, no sports event, no movie, no historical event that you can't record on one tape cassette with Omnivision IV.

Which means when a show you want to see runs longer than two hours (and don't the good ones always?) and you're miles away, or fast asleep, you don't have to worry about changing tapes, the way you'd have to with a two-hour system.

Okay, so you're convinced four hours of video taping is better than two hours. But there are a lot of other reasons to choose Omnivision IV VHS:

With our optional black and white video camera with sound, you can make your own TV shows. And they can be played on any VHS unit—regardless of brand.

You can use the same tape many times. Rerecording automatically erases old material. And to add to your video library of shows, sports and history-making events you've recorded, arrangements are now in progress for services that will allow you to transfer your slides and



When work rearranges your schedule, rearrange the network's schedule.

entire movie...even the climax when the Monster eats Yankee Stadium.

Four hours of taping when you are at home. So when the kids want to watch "Charlie Chocolate and His Merry Adventures" on one channel, you can still tape "The Monster that Ate New York" on another channel. And four hours means you can tape the en-

entire movie...even the climax when the Monster eats Yankee Stadium.

Four hours of video taping instead of two means there's virtually no program, no special, no sports event, no movie, no historical event that you can't record on one tape cassette with Omnivision IV.

Which means when a show you want to see runs longer than two hours (and don't the good ones always?) and you're miles away, or fast asleep, you don't have to worry about changing tapes, the way you'd have to with a two-hour system.

Okay, so you're convinced four hours of video taping is better than two hours. But there are a lot of other reasons to choose Omnivision IV VHS:

With our optional black and white video camera with sound, you can make your own TV shows. And they can be played on any VHS unit—regardless of brand.

You can use the same tape many times. Rerecording automatically erases old material. And to add to your video library of shows, sports and history-making events you've recorded, arrangements are now in progress for services that will allow you to transfer your slides and

home movies to VHS™ cassettes. Also prerecorded cassettes on a wide variety of subjects are available.

Omnivision IV™ can work with almost any TV set (although, naturally, we'd like to see it on a Panasonic TV... but that's another ad).

Omnivision IV VHS has Auto-Stop to automatically stop the tape when it runs out. Eliminates the need for you to be there.

To start tape recording when you're not at home, there's an electronic digital clock with a 24-hour timer.

And to find the exact place on the tape, there's memory rewind and a digital tape counter. To make recording convenient there's a remote pause control.

But that's not all: For long life and high performance there are HPF™ heads. Our annealed aluminum die-cast chassis means strength and stability. And a direct-drive motor provides greater picture stability than the belt-drive systems some of our competitors use.



Make a TV show with optional B/W camera and send it to Grandma or anyone with any VHS unit.

Can Omnivision IV VHS change your life? We think so. In fact, we think that the 4-hour video home system is the most important advance in television since television. But see it for yourself at selected dealers.



Build a video library of great shows, sports and historical events.




Catch up with your weekend chores while you catch the weekend game.



CAUTION: Unauthorized recording of copyrighted television programs, films, video tapes and other materials may infringe the rights of copyright owners and be contrary to copyright laws.

Panasonic **VHS**
Omnivision IV™

new books by
Bronowski
and the first
presidential
scientist



A Sense of the Future:
Essays in Natural Philosophy
By *J. Bronowski*

Published for the first time in book form, these essays by the author of *The Ascent of Man* reveal both the unity of his intellectual outlook and the diversity of his interests.

The major unifying theme of the book is the intensely creative and human nature of the scientific enterprise — its kinship with the highest levels of the artistic imagination and its ethical imperatives, which in guiding man's open-ended search for truth affirm his humanity.

Among the diversity of topics explored in the individual essays are the nature of human language in its biological context, intrinsic patterns of the mind, the limitations of logical systems, new concepts in evolution, humanism in the modern world, and the principle of tolerance.

\$12.50

Sputnik, Scientists, and Eisenhower:
A Memoir of the First Special Assistant to the President for Science and Technology
by *James R. Killian, Jr.*

The launching of Sputnik by the Russians 20 years ago created both blind panic and thoughtful reappraisals of American education. It also led Eisenhower to bring James Killian to Washington to redirect the thrust of the nation's scientific effort. Killian recounts those dramatic days and relates the lessons learned to today's issues.

\$14.95

The MIT Press

Massachusetts Institute of Technology
Cambridge, Massachusetts 02142




can begin anywhere, not just in regions of the highest reflectance, the first value in any sequence is arbitrarily assumed to be 100 percent. Because of this deliberately adopted fiction the sequential product becomes greater than unity whenever the path reaches an area whose reflectance is higher than that of the starting area.

The attainment of a sequential product greater than unity indicates that the sequence should be started afresh with the new area of high reflectance taken as being 100 percent. This procedure is the heart of the technique for finding the highest reflectance in the path. After the path reaches the highest reflectance in the scene, each of the sequential products computed thereafter becomes a fraction of the highest value. A satisfactory computer program has been designed to study the number of paths, their lengths and convolutions, the threshold values for recognizing edges and, perhaps most important, how to utilize all the pathways starting in all areas.

The biological counterpart of this

program is performed in undetermined parts of the pathway between the retina and the cortex. The process that corresponds to computing sequential products does not involve the averaging of areas or the averaging of flux. It does, however, call for an arithmetic that extends over the entire visual field. Furthermore, since the relevant phenomena are seen in a brief pulse of light, all the computations and conclusions about lightness must be carried out in a fraction of a second without dependence on eye movement. With a single pulse, eye movement, by definition, is not necessary. With continuous illumination the normal quick motions of the eye probably serve to maintain the freshness of the process.

With our computer model we can obtain a triplet of lightnesses for each area in the color Mondrian that corresponds closely to the lightnesses one would measure with a combined retinex filter and photomultiplier. The color corresponding to any given triplet can be visualized with the aid of the color "solid" we have built, in which the Munsell



SPOT OF LIGHT IN A VOID, that is, a single spot of narrow-band light viewed in an otherwise totally dark environment, has a color that would seem to depend solely on its wavelength. The color can also be explained, however, by the retinex theory in terms of lightness as perceived by the eye's three receptor systems. Psychophysical measurements show that when the eye is presented with a spot of light in a void, the perceived lightness is changed only slightly by very large changes in flux, as is indicated by the straight line. For example, if the spot is composed of a narrow-wavelength band centered, say, at 600 nanometers, the three cone pigments will absorb the flux in quite different amounts because of the shape of their absorption curves. In arbitrary units the long-wave pigment might absorb 80 units, the middle-wave pigment 20 units and the short-wave pigment a few tenths of a unit at most. If these ratios are plotted on the spot-in-a-void curve, the corresponding lightness values are 9 for the long waves, 8.5 for the middle waves and 7.5 for the short. This combination of lightnesses is perceived as a light reddish orange, not ordinarily seen under normal conditions unless surfaces are fluorescent.

colors are located in three dimensions in "lightness-color space" according to their lightness values measured in three wave bands through retinex filters [see illustration on page 120].

In normal images the sensation of white light will be generated by any area that is placed at the top of the lightness scale by all three retinex systems. On the other hand, an area that stands at the top of only two of the three lightness scales will be seen as some other color. Hence an area that is at the top of the lightness scale in the long- and middle-wave systems but is surpassed in lightness by some other area in the short-wave system will be seen not as white but as yellow. A similar intercomparison of triplets of lightnesses at the same place within each scene provides the sensation of color, area by area, in spite of unpredictable variations in illumination.

If one looks at black-and-white photographs taken through retinex filters, one sees a dramatic difference in lightness for most objects between the photograph representing the short-wave system and either of the photographs representing the other two systems. And yet it is the comparatively small differences between the long-wave and the middle-wave lightnesses that are responsible for the experience of vivid reds and greens.

Such reliable and sensitive responsiveness to small lightness differences provides the basis for the colors seen under anomalous conditions far removed from those the eye has evolved to see. Two examples of interest are the color of a spot of light in a total surrounding area devoid of light and the spectrum of colors produced by a prism.

One can readily measure the flux at the eye from a spot of light in a void. By changing the flux it is possible to estimate the corresponding change in perceived lightness. What one finds is that the estimated lightness changes only slowly with enormous changes in flux. For example, decreasing the flux by a very large amount will be seen as a very small reduction in lightness. If the spot of light is composed of a narrow band of long wavelength, say 600 nanometers, one can expect all three cone receptors to absorb the radiation in some degree, but significantly more radiation will be absorbed by the long-wave cones than by the other two kinds. When the three values are read on a scale of perceived lightness, the three lightnesses are 9 on the long-wave system, 8.5 on the middle-wave system and 7.5 on the short-wave system [see illustration on opposite page]. This combination of lightnesses is seen as a light reddish orange, a color not commonly perceived under ordinary conditions unless the surfaces are fluorescent. The spectrum, a strikingly anomalous display, can be regarded as a series of three laterally displaced

Fertilizing: Its importance in creating superior wines.



Fertilizing the vineyard is a vital part in the art of growing premium wine grapes. It is not a simple operation.

Its Importance

A healthy grapevine can thrive for 80 years or more. In fact, the older its stock, generally the better its grapes.

During the vine's lifetime its nutritional requirements will change from its earliest years—when it is being trained to the desired form for grape growing—to its later years—as it gradually matures into an established vine.

Thus, any fertilizer program that is to be complete must take into consideration both the vine's long-term needs and its immediate requirements.

Different Needs

Fertilizing is not a simple operation. We know that even within a single vineyard, there are no homogeneous soils. Soil textures can range from fine or coarse sand, to a very fertile sandy loam.

These different soils require different fertilizers in different amounts to compensate for deficiencies. As a result, we have kept a precise history of all the fertilizing experiences in our vineyards over the past 30 years.

We know exactly when, how much, and what kind of fertilizer has been applied to each of our vineyards during that time.

We know the results of those applications—and how long the treatment lasted—down to small problem areas only a few yards square and even individual vines.

One of the methods we use to gauge these results is core sampling—the extraction of a cross-section of earth to be analyzed for nutrients—or the lack of them. Above, you see a rendition of our core sampler being used.

We take samples at various depths down to four feet from any part of the vineyard which may show abnormal vine growth or visual deficiencies for comparison with samples from normal areas. These soil samples are delivered to our winery for immediate analysis. They will be analyzed and compared for available nitrogen, pH, organic matter, and other important elements in order to de-

termine the source of the problem and the type and quantity of fertilizer which is required.

Fertilizing As An Exact Science

There are two good reasons why we take so much care when it comes to fertilizing: under-fertilizing and over-fertilizing. Under-fertilizing greatly reduces vine growth and bud fruitfulness, whereas over-fertilizing can produce such exceptionally heavy leaf growth as to curtail bud fruitfulness and grape production.

These phenomena are the result of an imbalance of what is known as the carbohydrate-nitrogen level of the vine.

A high carbohydrate-moderate nitrogen level produces moderate vegetative growth, early wood maturity, abundant fruit bud formation and good fruit production.

This is what we look for in the Gallo Vineyards.

Petiole Analysis

In addition to normal soil analysis in our vineyards we also employ a test known as "petiole analysis."

The petiole is the stalk of a vine leaf. By taking a sample of it and analyzing it, we can determine the current nutritional status of the entire grapevine.

If we find that a particular area of one of our vineyards is weaker than the rest, we will immediately gather 200 or more petiole samples from that section—plus a like number from a healthier area of the same vineyard for comparison.

By analyzing the tissues of all these samples, we can readily and specifically determine if the weaker area has a deficiency of nitrogen, potassium, zinc, boron or some other vital mineral.

Our Solution

These are only a few of the many special steps we take in the Gallo Vineyards to try to grow the finest grapes possible.

It is only by such meticulous care along the entire chain of grape growing and winemaking steps that we can hope to achieve our goal:

To provide you with the finest wines we can produce.

Gallo Vineyards, Modesto, California

HP measurement and computer advances

The HP-67. The most powerful hand-held programmable HP has ever built. Handles programs of up to 224 steps on a single magnetic card while larger programs—even thousands of steps—are easily handled with several cards. Automatic merging of keycodes increases the HP-67's program capacity. \$450*.



The HP-97. All the capability of the HP-67, plus a built-in thermal printer. Small enough to fit very portably into your briefcase. \$750*.

The HP-67/HP-97 software library: turnkey solutions to your computational problems

...whether you work in astronomy, surveying, computer science, accounting, thermodynamics, biology, probability, medical science, number theory, machine design, real estate, aviation, marine navigation, solid state devices, behavioral science, dynamics, differential equations, quality assurance, loans, optics, statistics, (or you name it).

We even have a program that converts your calculator into a stopwatch. And one that computes your biorhythm. There is one that tells time from the length of your shadow. And one that solves the classic coconut problem.

Or you can create your own problem solver. With the clear instructions provided in the Owner's Handbook, it is so easy you'll be writing programs the first day. And once you've keyed in a program you can permanently record it on a magnetic card.

Here's how to build your personal program library:



- With your new HP-67 or HP-97 you'll receive a Standard Application Pac containing 15 programs from a variety of fields. These are pre-recorded on magnetic cards, which means that in less than two seconds your calculator will be ready to solve complex problems at the touch of a few keys.
- Choose from 10 application pacs each containing an instruction manual and up to 25 pre-



corded programs in specific fields such as math, electrical and mechanical engineering, statistics, and surveying. Applications pacs are \$35* each.

- Choose from more than 1000 programs in the HP-67/HP-97 Users' Library, containing programs contributed by HP-67 and HP-97 owners as well as by Hewlett-Packard engineers. These programs are not pre-recorded, but consist of step-by-step instructions for keying them into your calculator. They're only \$3* each.



- Choose from 40 new Users'-Library Solution Books. Each is a collection of 10 to 15 Users' Library programs in fields ranging from aeronautical engineering to thermal and transport sciences. Solutions Books are \$10* each.

If you're curious about the range of HP Users' Library programs, send us this coupon. We'll send you a free catalog listing all the contributed programs as well as the contents of each Solutions Book. Our working combination of powerful calculators and comprehensive programs can deliver accurate solutions to your problems—painlessly.

extend your possibilities.



For those who design or debug digital system hardware and software, this new logic analyzer is a major breakthrough.

The new HP 1615 is the first logic analyzer that can make state and timing measurements simultaneously and relate one to the other. It is also the first that can detect timing glitches, strip them from captured data, and use them to trigger further analyses.

Digital machines operate both synchronously and asynchronously. A machine's program is implemented in synchronous mode by system clocks that change the state of the machine. Logic state analyzers—introduced by Hewlett-Packard about four years ago—do a great job of measuring synchronous activity: they monitor program flow and tell the designer when machine execution deviates from the program.

But much of the activity of a digital system is asynchronous—as when the program calls for something to happen between clock pulses, e.g. to move data on an I/O structure or through an address decoder. If such an activity is not carried out on the specified time schedule, the system malfunctions. Timing analyzers are used to measure asynchronous activity.

Even with separate state and timing measurements in hand, the designer does not have a really good view of overall system activity until he can relate any asynchronous fault to the state of the machine at the time of the fault. And that's been an extremely tricky proposition, one that has been left to the designer's ingenuity and intuition.

The HP 1615 performs both state and timing analyses *simultaneously* and *interactively*. Because of this dual capability, it can capture any asynchronous activity that occurs at a specific point in the program; identify which program step a system was executing at the time of malfunction; or monitor handshake sequences at specific points in program execution. Before the introduction of the HP 1615, no single instrument had been able to make these objective measurements.

Then there's the matter of glitches, those digital gremlins that confuse the machine's logic. Because they are transient and unpredictable, problem-causing glitches are extremely difficult to distinguish from harmless ones. Now the HP 1615 does it all: (1) captures system activity, both synchronous and asynchronous, simultaneously, (2) captures related data on surrounding lines, (3) strips the glitches out of this data, and (4) uses the glitches as part of the trigger for state, timing, or combined state and timing analyses.

The HP 1615 can be configured, through simple keyboard entries, as a 24-bit state analyzer, or as an 8-bit timing analyzer, or as a combined 16-bit state and 8-bit timing analyzer. It can trace events in as many as 24 channels at rates to 20 MHz, capturing data in a memory that is 24 bits wide and 256 words deep.

Although most system designers and debuggers will declare the HP 1615 to be priceless, it can be purchased for \$6800*.



1503 Page Mill Road, Palo Alto, California 94304

For assistance call: Washington (301) 948-6370, Chicago (312) 255-9800, Atlanta (404) 955-1500, Los Angeles (213) 877-1282.

Mail to: Hewlett-Packard, 1503 Page Mill Road, Palo Alto, CA 94304
Please send me further information on

- HP-67/HP-97 software library
- HP 1615 logic analyzer

Name _____
Company _____
Address _____
City _____ State _____ Zip _____

*Domestic U.S. prices only

00840

continuous gradients involving both the properties of spots and the properties of areas. From these properties it is possible to predict the colors of the spectrum, whereas it is not possible, as we have seen, to attribute a specific spectral composition to the radiance from a colored area in everyday life.


Perhaps the first observation pointedly relevant to the mechanism of color formation in images is not Newton's spectrum but the phenomenon of colored shadows, described in 1672 by Otto von Guericke. "This is how it happens," he wrote, "that in the early morning twilight a clear blue shadow can be produced upon a white piece of paper [by holding] a finger or other object... between a lighted candle and the paper beneath." This important experiment, we now know, depicts an elementary example of generating three different lightnesses on the three receptor systems. A diagram of this experiment with long-wave ("red") light and white light appears below. Here the color of the shadow is blue-green. The diagram shows that the triplet of lightnesses in the shadow corresponds to the blue-green color

one would predict for it from its position in lightness-color space.

One can now understand the red-and-white images of our early work as a procedure that carries the colored shadow to a richly variegated family of colors no longer in shadows but in images. The colors seen in a red-and-white projection can be readily predicted by extending the analysis followed in predicting the color of von Guericke's shadow. To demonstrate this point we reproduce on page 128 the "red" and "green" separation images used in making a red-and-white multicolored projection. (In this demonstration no attempt is made to reproduce the colors seen in the actual multicolored image.) The red-and-white projection was photographed through long-, middle- and short-wave retinex film-filter combinations. The three images are reproduced below the pair of long- and middle-wave separation images that were superposed to make the red-and-white image. The significant point is that when the eye views the red-and-white images on the screen with its own retinex system, it is provided with a triplet of lightnesses for each part of the scene that resembles the triplet it would

obtain if it viewed the original scene directly. In this important meeting point of the blue-green shadows with the colored images, provided by the red-and-white display, the extended taking and multiplication of ratios determine the lightness of each small area. Finally, all these principles are applied in everyday ternary vision, which creates a distinct lightness image for each of the three sensitive systems and compares them in order to generate color.

The train of interlocking concepts and experiments started 25 years ago with the observation that the relative energies of the red-and-white projectors can be altered without changing the names of the various colors. This observation negated the simplistic explanation in terms of contrast, fatigue and surround and led to the fundamental concept of independent long- and short-wave image-forming systems that ultimately evolved to the concept of three independent retinex systems and to the Mondrian demonstration. The concept of the percentage of available light on each wave band as a determining variable and the technique of measuring it



BLUE-GREEN COLORED SHADOW is seen when a hand or some other object is placed in the beam of a projector that is sending long-wave ("red") light to a screen while the screen is illuminated by a beam of white light. The author regards Otto von Guericke's description in 1672 of seeing colored shadows made by candlelight as the first observation pointedly relevant to the mechanism of image and color formation. In the analysis at the right it is assumed that one projector sends white light to the screen. The other projector, equipped with a red filter, sends only long wavelengths to the screen. Assume that the white light contributes 100 arbitrary units of flux to each of the short-, middle- and long-wave receptors. The long-wave flux is absorbed by the three receptor systems in different proportions: 100 units are

absorbed by the long-wave system, 50 by the middle-wave system and five by the short-wave system. (A small amount of scattered long-wave light also appears in the shadow.) The third column of boxes shows the combined amounts of flux from both sources absorbed by each receptor system. The fractions represent the ratio at edges of the flux from within the shadow divided by the flux from outside. The fourth column shows the lightness on each receptor system. The lightness of the lightest place in the scene for each receptor system will be near the top of the lightness scale, being determined by the flux of radiant energy in the same way that a spot has its lightness determined by flux. Triplet of lightnesses within the shadow falls in the region of color space that the eye perceives as being blue-green.

MORE gears—with 5-speed transmission standard or optional automatic.*

MORE rubber—with wider 185 steel belted radials.

MORE stability—with a front end spoiler.

MORE interior selections—with tartan plaid or brushed solid nylon broadcloth.

MORE control—with a lower suspension.

AND STILL MORE:

2-liter overhead cam engine.

Rack and pinion steering.

Full instrumentation.

MacPherson strut

front suspension.

Power-assisted front disc brakes. A cockpit wider than a Corvette or a Z-Car's.

Triumph TR7. The shape of things to come.

Now with more.




TRIUMPH

For the name of your nearest Triumph dealer call: 800-447-4700. In Illinois call: 800-322-4400. British Leyland Motors Inc., Leonia, N.J. 07605.

*Automatic transmission not available in California.

MORE TR7



THE SHAPE OF THINGS TO COME
TR7


evolved to the concept that lightnesses maintain an independent rank order on long- and short-wave bands. This measuring technique in turn evolved from a projected black-and-white image to an arrangement of colored papers in the color Mondrian. The manifest stability and constancy of the lightnesses of all the papers of the Mondrian when a single wave band illuminates it with varying intensity dramatizes the concept that every colored paper has three reflectances on three wave bands and that these reflectances are somehow connected with the biological characteristic: lightnesses.

A black-and-white Mondrian taught that nonuniformity of illumination, size and shape of area and length of edges were basically irrelevant to light-

ness. What was needed was a far-reaching, edge-reading arithmetic: the sequential product of ratios at edges. For the color Mondrian the ratio at edges was early recognized as requiring a ratio of the integrals of the product at each wavelength of the absorbance of the cone pigment times the reflectance of the colored paper times the illuminants. Separate integrals were taken over the wave bands of the three cone pigments. In a long series of binocular comparison-and-selection observations the quantity satisfying the integral was shown to be impressively well correlated with lightness, particularly after the realization that the scale, or spacing, of the reflectance integral should be made to correspond with the spacing of the biological quantity lightness. This led to the designation "scaled integrated re-

flectance" as the external partner to which the retinex system relates the internal partner: constructed lightness.

Color can be arranged in the lightness solid with long-, middle- and short-wave axes of lightness. All visible colors reside in this solid independent of flux, each color having a unique position given by the three axial values of lightness. It should be remembered that the reality of color lies in this solid. When the color Mondrian is nonuniformly illuminated, photographed and measured, reflectance in the photograph no longer correlates with the color but the lightness does. The three sets of ratios of integrals at edges and the product of these integrals within a set emerge as the physical determinants in the partnership between the biological system and areas in the external world.



RETINEX RECORDS OF RED-AND-WHITE projections show that red-and-white images produce a triplet of lightnesses for each part of the scene that are consistent with the observed color sensations. The two photographs in the top half of this illustration are reproductions of the long-wave (*left*) and middle-wave (*right*) separation records taken of the original still life. The long-wave record was projected onto a screen with a long-wave (red) filter in the beam of light. The middle-wave record was projected in superposition onto the same screen in the light of a tungsten-filament lamp. Three retinex photographs were then taken of projected images appearing on

screen. The retinex records are reproduced in the bottom part of the illustration: long-wave at the left, middle-wave in the middle and short-wave at the right. The colors seen in red-and-white projections are those one would expect from their triplets of lightnesses. The apple is light on the long record and darker in the middle and short records. The orange is lightest on the long record, intermediate on the middle record and darkest on the short. It is impressive that with his own retinex systems the observer can see a blue cup, a brown straw basket and pale yellow bananas with lightness differences so small as to challenge photoengraving process used to reproduce photographs.